

RESULTS AND RECOMMENDATIONS OF THE 7TH EUROPEAN HYPOXIA SYMPOSIUM

Markus Tannheimer¹, Corado Angelini^{2,3}, Martin Faulhaber⁴, David Hillebrandt⁵, Zbigniew Szygula⁶, Hans-Volkhard Ulmer⁷, Anton Usaj⁸, Rianne van der Spek^{3,9}, Thomas Küpper^{3,10}

¹Department of Sport and Rehabilitation Medicine, University of Ulm, Ulm, Germany

²Department Neurosciences University of Padova and IRCCS Fondazione San Camillo Hospital of Venice, Venice, Italy

³Medical Commission of the Union Internationale des Associations d'Alpinisme (UIAA MedCom), Bern, Switzerland

⁴Department of Sport Science, University Innsbruck, Innsbruck, Austria

⁵President UIAA Medcom, Vice President ISMM, Hon Medical Advisor to British Mountaineering Council

⁶Department of Sports Medicine and Human Nutrition, Faculty of Physical Education and Sport, University of Physical Education, Krakow, Poland

⁷Institute of Sport Science, University of Mainz, Mainz, Germany

⁸Laboratory of Biodynamics, Faculty of Sport, University of Ljubljana, Ljubljana, Slovenia

⁹Department of Endocrinology & Metabolism, Academic Medical Center - University of Amsterdam, Amsterdam, The Netherlands

¹⁰Department of Occupational and Social Medicine, RWTH Aachen University, Aachen, Germany

One of the main themes of the 7th European Hypoxia Symposium (<http://www.hypoxiasymposium.org>; September 19-21st. 2014, Berchtesgaden, Germany) was „Contraindications for exposure to Hypoxia“ [1]. Participants from 10 countries engaged in an animated discussion with the speakers that presented data on various aspects of high altitude sojourns with pre-existing conditions. In this paper, we summarize the statements and recommendations that resulted from the presentations and discussions.

Exposure to real or artificial altitude (normobaric hypoxia) is increasingly common [2-4], not only in classical mountaineering but in a growing number of occupational settings as well. For example, hypoxia is used to protect cultural heritage, and scarcity of natural resources leads to mining at higher altitudes. In public opinion, climbing and mountaineering are often perceived as dangerous [5]. However, objective data grade mountaineering and climbing as relatively harmless [6]. Trekking is associated with a minor risk of accidents and it is more and more popular amongst elderly people (mean 42.7 yrs.: min 18, max 76 yrs.) [7]. Therefore the number of trekkers with pre-existing medical conditions is rising [8], as is the number of physicians' consultations prior to a planned high altitude sojourn. High altitude medicine and alpine medicine are rarely taught in the medical university system [9], and as a result most consulted physicians

tend to be too cautious in their recommendations [2]. To support doctors that are consulted in hypoxia or high altitude related matters, the Medical Commission of the UIAA (Union Internationale des Associations d'Alpinisme) has published 24 recommendations. The topics of these recommendations range from travelling at altitude with pre-existing cardiovascular conditions to the use of portable hyperbaric chambers, and they are available in several languages, including German (http://www.theuiaa.org/medical_advice.html).

Anyone travelling to high altitude should be aware of high altitude diseases and should know what to do when symptoms develop. Good sources of information include www.ismmmed.org and www.medex.org.uk [10].

Risk assessment in clinical practice should start with evaluation of any cardiopulmonary diseases that could worsen during a sojourn involving high altitude [10]. For persons with no previous experience of prolonged stay at high altitude, risk assessment is difficult [11,12]. The most reliable predicting factor for developing high altitude related disease is previous episodes in the patients' medical history [10]. If previous sojourns were unproblematic, further sojourns with comparable rates of ascent and final altitude will be most likely be unproblematic too [13]. Mountaineers should therefore be advised to initially choose a tour with moderate demands. The rate of

ascend is of crucial importance [14], and a simple altitude profile graph provides clear insight into the risk of developing altitude related diseases. Above 2500 m acclimatization is necessary but time needed for proper acclimatization varies between individuals [15]. Therefore, the daily gain in sleeping altitude should not exceed 300–500 m [16]. Furthermore, after every 1000 m of altitude gain an additional day for proper acclimatization is recommended [17]. These easy recommendations are frequently ignored [18]. For example, the incidence of acute mountain sickness (AMS) on Mount Kilimanjaro was reported to range between 47% and 75% [19-21], and high altitude pulmonary edema (HAPE) occurs often [22]. Despite the relative lack of technical difficulties, only 53% of climbers reach the summit [19]. The high incidence of avoidable high altitude related diseases make this peak one of the most dangerous mountains in the world [23].

When high altitude disease does occur, or pre-existing medical conditions worsen, the aetiology is of major interest. In many cases altitude profiles were too ambitious, a frequent problem in commercially offered tours [18,24]. Common causes are group dynamics, burning ambition or even lack of knowledge among the participants, as well as economic motives from the company side. As early as 1992 Shlim [25] showed that amongst trekkers in Nepal, mortality was 5-fold higher in commercial tours compared to individually organized tours. According to the preliminary data of the ADEMED-Expedition (www.ademed.de) [7], nothing has changed up to now.

In the advisory setting, the fundamental question is: 'will the pre-existing medical condition aggravate during the high altitude stay or can it compromise the safety of the patient respectively his partner?' [2,26]. For example, a hypoglycemic episode in a patient with type 1 diabetes endangers both the patient and his climbing partner. From an ethical point of view, it is necessary to inform the climbing partner about relevant diseases. Returning to our diabetic patient, in case of well-trained handling of the disease, even multi-pitch alpine climbing tours should be possible [27]. The UIAA recommendation 'Drugs at altitude': gives a good overview of temperature stability of many frequently used drugs [28].

Conclusions and Consensus

In general, even for people with pre-existing medical conditions, high altitude sojourns should be possible, provided fundamental recommendations of acclimatization are followed [11]. The UIAA guidelines have been developed to assist physicians without high altitude expertise to give advice in uncomplicated cases. In complex cases or severe pre-existing medical conditions, advice of a physician with expertise in

high altitude (and travel) medicine should be sought to assess specific individual risks [8,26].

Declaration of interest

The authors report no conflicts of interest.

References

1. Ulmer HV. Medical Contraindications for Mountaineering or Flying in Hypoxia - Introductory Considerations. *Med Sport* 2014; 18(3): 122.
2. Küpper T. Hypoxia and Cardiac Diseases - where are the Limits? *Med Sport* 2014; 18(3): 115.
3. Lechner R. Military Missions at High Altitude - tactical Aspects of High Altitude Physiology. *Med Sport* 2014; 18(3): 116.
4. Tannheimer M. Craniocerebral Injury during Expedition Climbing. *Med Sport* 2014; 18(3): 120.
5. Becher S. Mountain Sportive Activity - too Dangerous for Private Insurers? *Med Sport* 2014; 18(3): 109.
6. Küpper T, Kühn J, Scheider M, et al. Risk in Mountain Sports - Facts and Data. *Med Sport* 2014; 18(3): 115.
7. Müller-Ost M, Schmitz S, Apel C, et al. Cardiovascular Risk profiles and pre-existing Conditions in Trekkers - an Epidemiological Study in the Solo Khumbu Region, Nepal. Data from the 2nd ADEMED Expedition 2011. *Med Sport* 2014; 18(3): 117-8.
8. Rimoldi SF, Sartori C, Seiler C, et al. High-altitude exposure in patients with cardiovascular disease: risk assessment and practical recommendations. *Prog Cardiovasc Dis* 2010; 52(6): 512-24.
9. Treff G, Tannheimer M, Sareban M, et al. A practical Approach to Altitude Medicine for Students of Sports Science and Medicine. *Med Sport* 2014; 18(3): 121.
10. Bartsch P, Swenson ER. Clinical practice: Acute high-altitude illnesses. *N Engl J Med* 2013; 368(24): 2294-302.
11. Richalet JP, Canoui-Poitine F. Pro: hypoxic cardiopulmonary exercise testing identifies subjects at risk for severe high altitude illnesses. *High Alt Med Biol* 2014; 15(3): 315-7.
12. Faulhaber M, Pocecco E, Gatterer H, et al. Resting Arterial Oxygen Saturation for the Prediction of subsequent Development of Acute Mountain Sickness. *Med Sport* 2014; 18(3): 112.
13. Bartsch P. Con: hypoxic cardiopulmonary exercise testing identifies subjects at risk for severe high altitude illnesses. *High Alt Med Biol* 2014; 15(3): 318-20.
14. Luks AM, McIntosh SE, Grissom CK, et al. Wilderness Medical Society consensus guidelines for the prevention and treatment of acute altitude illness. *Wilderness Environ Med* 2010; 21(2): 146-55.
15. Tannheimer M, Albertini N, Ulmer HV, et al. Testing individual risk of acute mountain sickness at greater altitudes. *Mil Med* 2009; 174(4): 363-9.
16. Luks AM, McIntosh SE, Grissom CK, et al. Wilderness Medical Society consensus guidelines for the prevention and treatment of acute altitude illness. *Wilderness Environ Med* 2010; 21(2): 146-55.
17. Kubalova J. 4 x 4 Health Rules for Mountaineers. Consensus Statement of the UIAA Medical Commission: Advice and Recommendations 2008 http://theuiaa.org/medical_advice.html
18. Shah NM, Windsor JS, Meijer H, et al. Are UK commercial expeditions complying with wilderness medical society guidelines on ascent rates to altitude? *Journal of Travel Medicine* 2011; 18(3): 214-6.
19. Karinen H, Peltonen J, Tikkanen H. Prevalence of acute mountain sickness among Finnish trekkers on Mount Kilimanjaro, Tanzania: an observational study. *High Altitude Medicine & Biology* 2008; 9(4): 301-6.
20. Jackson SJ, Varley J, Sellers C, et al. Incidence and predictors of acute mountain sickness among trekkers on Mount Kilimanjaro. *High Altitude Medicine & Biology* 2010; 11(3): 217-22.

21. Eigenberger P, Faino A, Maltzahn J, et al. A Retrospective Study of Acute Mountain Sickness on Mt. Kilimanjaro Using Trekking Company Data. *Aviation, Space, and Environmental Medicine* 2014; 85(11): 1125-9.
22. Cohn V. Tennis Star's Hospitalization for Altitude Sickness During Mount Kilimanjaro Ascent Highlights Risks of High Altitude Climbs. 2010 <http://www.liebertpub.com/global/pressrelease/tennis-stars-hospitalization-for-altitude-sickness-during-mount-kilimanjaro-ascent-highlights-risks-of-high-altitude-climbs/848/>
23. Windsor JS. Safety and Success on Kilimanjaro. Consensus Statement of the UIAA Medical Commission: Advice and Recommendations 2008 http://www.theuiaa.org/upload_area/files/1/Kilimanjaro-UIAA%281%29.pdf
24. Tannheimer M. Intermittent simulated hypoxia for pre-acclimatization. *Sleep Breath* 2010; 14(3): 185-6.
25. Shlim DR, Gallie J. The causes of death among trekkers in Nepal. *Int J Sports Med* 1992; 13 Suppl 1: S74-6.
26. Angelini C. Contraindications in Neurological Disorders in the Mountains. *Med Sport* 2014; 18(3): 108.
27. Hillebrandt D. Diabetes: Personal practical Experience of Management of Diabetes in the Field. *Med Sport* 2013; 17(3): 105.
28. Küpper T, Milledge J, Basnyat B, et al. The Effect of Extremes of Temperature on Drugs. 2008 http://theuiaa.org/upload_area/Medicine/Advice-and-Recommendations/No10-Drugs/UIAA_MedCom_Rec_No_10_Drugs_at_Altitude_2008_V1-2.pdf

Accepted: November 10, 2014

Published: November 14, 2014

Address for correspondence:

Markus Tannheimer

Department of Sport and Rehabilitation Medicine

University of Ulm

Leimgrubenweg 12-14

89070 Ulm, Germany

Markus.Tannheimer@arcor.de

Corrado Angelini: corrado.angelini@unipd.itMartin Faulhaber: martin.faulhaber@uibk.ac.atDavid Hillebrandt: dh@hillebrandt.org.ukZbigniew Szygula: wfszygul@cyf-kr.edu.plHans-Volkhart Ulmer: ulmer@uni-mainz.deAnton Usaj: anton.usaj@fsp.uni-lj.siRianne van der Spek: riannevdspek@gmail.comThomas Küpper: tkuepper@ukaachen.de**Authors' contribution**

A – Study Design

B – Data Collection

C – Statistical Analysis

D – Data Interpretation

E – Manuscript Preparation

F – Literature Search

G – Funds Collection